

# Flebogamma® 5% DIF

Immune Globulin Intravenous  
(Human)

Pure confidence

## Infusion Rate Calculations in mL (cc) per Hour

Patient's Weight		kg	10	20	30	40	50	60	70	80	90	100	110	120
		lb	22	44	66	88	110	132	154	176	198	220	242	264
Infusion Rate														
mL/kg/min	mL/kg/hr													
0.01	0.6		6	12	18	24	30	36	42	48	54	60	66	72
0.02	1.2		12	24	36	48	60	72	84	96	108	120	132	144
0.03	1.8		18	36	54	72	90	108	126	144	162	180	198	216
0.04	2.4		24	48	72	96	120	144	168	192	216	240	264	288
0.05	3.0		30	60	90	120	150	180	210	240	270	300	330	360
0.06	3.6		36	72	108	144	180	216	252	288	324	360	396	432
0.07	4.2		42	84	126	168	210	252	294	336	378	420	462	504
0.08	4.8		48	96	144	192	240	288	336	384	432	480	528	576
0.09	5.4		54	108	162	216	270	324	378	432	486	540	594	648
0.10	6.0		60	120	180	240	300	360	420	480	540	600	660	720

### Shaping the future. See the difference today

The infusion of Flebogamma® 5% DIF should be initiated at a rate of 0.01 mL/kg body weight/minute (0.5 mg/kg/minute). If, during the first 30 minutes, the patient does not experience any discomfort, the rate may be gradually increased to a maximum of 0.10 mL/kg/minute (5 mg/kg/minute).